

Galeno - Modeling and Control for Personalized Drug Administration

Standard PK/PD Model

Rocuronium

This work was supported by FEDER founds through *COMPETE*-Operational Programme Factors of Competitiveness ("Programa Operacional Factores de Competitividade") and by Portuguese founds through the *Center for Research and Development in Mathematics and Applications* (University of Aveiro), the Portuguese Foundation for Science and Technology ("FCT-Fundação para a Ciência e a Tecnologia"), within project PEst-C/MAT/UI4106/2011 with COMPETE number FCOMP-01-0124-FEDER-022690 and the Project GALENO – Modeling and Control for Personalized Drug Administration, FCT PTDC/SAU-BEB/103667/2008.

2012, October



Standard PK/PD Model - Rocuronium

The most common models for the effect concentration of a drug are compartmental systems.

A system is a set of interconnected elements that are dependent on each other and form a unit that has specific characteristics and functions. Each system admits states that are defined as a set of variables capable to describe the system in any instant of time. Systems can be classified as continuous or discrete-time systems according to the continuous or discrete nature of the time-line over which their variables are defined. Also, systems can be classified as open, closed or isolated. An open system can exchange matter or energy with its surroundings, while a closed system can only exchange energy, but not matter with its surroundings. In contrast, an isolated system cannot exchange neither energy nor matter.

Compartmental systems are widely used to model the pharmacodynamics and pharmacokinetics of intravenously administered drugs. A compartmental system is a system that has a finite number of homogeneous, well-mixed subsystems, called compartments that exchange material among them and with the environment. These models are based on the principle of mass conservation.



Figure 1 - Representation of a compartment.

Fig. 1 represents a compartment, in this case compartment i; here b_i represents the input rate (the drug infusion rate to a patient if we consider the specific case of anesthesia), x_i is the concentration of material in the compartment i, k_{ij} represents the rate of mass transfer from compartment i to compartment j, and k_{i0} represents the rate of material output from compartment i to the environment. The input to compartment i, is given by $b_i.u$, where u is the total system input. This input, the state x_i and all the rate constants are assumed to be non-negative. At each time instant t, the variation $\dot{x}(t)$ is the concentration of material in compartment i.



Figure 2 - Wiener Model of the Standard PK/PD Model for Rocuronium



Compartmental Model (Linear Part)

Here this model includes three compartments, where two of them represent the pharmacokinetic model (central and peripheral compartments), combined with an effect compartment which is part of the pharmacodynamic part of the model, see Fig. 3. Pharmacokinetics describes the path that a drug does inside the body, whereas pharmacodynamics means the pharmacology field that studies the physiological effects of drugs, and their mechanisms of action.



Figure 3 - Compartmental Model of the Standard PK/PD Model for Rocuronium.

This model has six patient dependent parameters, k_{10} , k_{12} , k_{1e} , k_{20} , k_{21} , and k_{e0} (min⁻¹) that must be identified for each particular patient. The state variables x_1 and x_2 (µg/kg) correspond to the drug concentrations in central and peripheral compartments, whereas x_3 (µg/kg) is the drug concentration in the effect compartment, also known as effect concentration. The input u(t) (µg/kg/min) corresponds to the delivery rate of drug concentration with respect to the central compartment, and is computed as $u(t) = \tilde{u}(t)/V_1$, where $\tilde{u}(t)$ is the drug delivery rate, and V_1 is the volume of the central compartment. The output y(t) (µg/kg) corresponds to the effect concentration.

The corresponding state-space equations are as follows.

$$\begin{bmatrix} \dot{x_1} \\ \dot{x_2} \\ \dot{x_3} \\ \dot{x} \end{bmatrix} (t) = \begin{bmatrix} -(k_{12} + k_{10} + k_{1e}) & k_{21} & 0 \\ k_{12} & -(k_{21} + k_{20}) & 0 \\ k_{1e} & 0 & -k_{e0} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \dot{x} \end{bmatrix} (t) + \begin{bmatrix} 1 \\ 0 \\ 0 \\ Matriz B \end{bmatrix} u(t)$$

$$y(t) = \underbrace{[0 \ 0 \ 1]}_{Matriz C} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \dot{x} \end{bmatrix} (t)$$



Hill Equation (Nonlinear Part)

As an output from the models presented in the previous subsection, the effect concentration, y(t), is obtained. The effect concentration corresponds to the percentage of administered drug that will produce effect in the NMB level, r(t) (%). The relationship between the effect concentration and the NMB level is given by the Hill Equation; this is a nonlinear static equation.

$$r(t) = \frac{100}{1 + \left(\frac{y(t)}{C_{50}}\right)^{\gamma}},$$

where C_{50} (µg/kg) and γ (gamma) (dimensionless) are patient-dependent parameters.

The equations obtained from the state-space equations are the following:

- $\frac{dx_1}{dt} = u(t) x_1 \cdot (k_{12} + k_{10} + k_{1e}) + x_2 \cdot k_{21}$
- $\frac{dx_2}{dt} = x_1 \cdot k_{12} x_2 \cdot (k_{21} + k_{20})$

•
$$\frac{dx_3}{dt} = x_1 \cdot k_{1e} - x_3 \cdot k_{e0}$$

Since:

•
$$V_i c_i = x_i$$

•
$$\frac{dC_i}{dt} = \frac{1}{V_i} \times \frac{dx_i}{dt}$$

where V_i = 1:

•
$$\frac{dC_1}{dt} = u(t) - (k_{12} + k_{10} + k_{1e}) + k_{21}$$

• $\frac{dC_2}{dt} = k_{12} - (k_{21} + k_{20})$
• $\frac{dC_3}{dt} = k_{1e} - k_{e0}$

Effect Concentration (C_e) - $\frac{dC_e}{dt} = \frac{dC_3}{dt}$

Bibliography

[1] - E. P. MacCoy, R. K. Mirakhur, V. R. Maddineni, J. M. K. H. Wierda and J. H. Proost, "Pharmacokinetics of Rocuronium after Bolus and Continuous Infusion during Halothane Anaesthesia", British Journal of Anaesthesia, 1996, 76, 29-33.