

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Editors: **W. Clegg and D. G. Watson**

2,6-Diphenylpyridine

Marisa A. Rocha, John Nicolson Low, Ligia R. Gomes, Antonio Quesada and Luis M. N. B. F. Santos

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Reproduction of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>

2,6-Diphenylpyridine

Marisa A. Rocha,^a John Nicolson Low,^{b*} Ligia R. Gomes,^c
Antonio Quesada^d and Luís M. N. B. F. Santos^a

^aCentro de Investigação em Química, Departamento de Química, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre 687, P-4169-007 Porto, Portugal, ^bDepartment of Chemistry, University of Aberdeen, Meston Walk, Old Aberdeen AB24 3UE, Scotland, ^cCIAGEB-Faculdade de Ciências da Saúde, Escola Superior de Saúde da UFP, Universidade Fernando Pessoa, Rua Carlos Maia 296, P-4200-150 Porto, Portugal, and ^dDepartamento de Didáctica de las Ciencias, Universidad de Jaén, 23071 Jaén, Spain
Correspondence e-mail: che562@abdn.ac.uk


Received 7 November 2007; accepted 13 November 2007

Key indicators: single-crystal X-ray study; $T = 120\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.055; wR factor = 0.163; data-to-parameter ratio = 9.8.

In the title compound, $C_{17}\text{H}_{13}\text{N}$, the dihedral angles between the pyridine ring and the phenyl rings are $29.68(18)$ and $26.58(17)^\circ$. In the crystal structure, the molecules are linked by a weak $\text{C}-\text{H}\cdots\pi$ interaction, leading to $[0\bar{1}1]$ chains. There are no further significant intermolecular interactions.

Related literature

For related literature, see: Crispini & Neve (2002); Silva *et al.* (1997). For a previous synthesis, see: Miyaura & Suzuki (1995).



Experimental

Crystal data

$C_{17}\text{H}_{13}\text{N}$	$V = 1273.9(2)\text{ \AA}^3$
$M_r = 231.28$	$Z = 4$
Orthorhombic, $Pna2_1$	Mo $K\alpha$ radiation
$a = 16.1368(16)\text{ \AA}$	$\mu = 0.07\text{ mm}^{-1}$
$b = 12.5371(14)\text{ \AA}$	$T = 120(2)\text{ K}$
$c = 6.2969(4)\text{ \AA}$	$0.70 \times 0.32 \times 0.14\text{ mm}$

Data collection

Bruker-Nonius KappaCCD diffractometer	29987 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 2003)	1596 independent reflections
$T_{\min} = 0.953$, $T_{\max} = 0.990$	1147 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.053$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$	1 restraint
$wR(F^2) = 0.163$	H-atom parameters constrained
$S = 1.07$	$\Delta\rho_{\max} = 0.21\text{ e \AA}^{-3}$
1596 reflections	$\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$
163 parameters	

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg1$ is the centroid of the pyridine ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$C64-\text{H}64\cdots Cg1^i$	0.95	2.74	3.534 (4)	142

Symmetry code: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *COLLECT* (Nonius, 1999); cell refinement: *DIRAX/LSQ* (Duisenberg *et al.*, 2000); data reduction: *EVALCCD* (Duisenberg *et al.*, 2003); program(s) used to solve structure: *FLIPPER* (Oszlányi & Sütő, 2004, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97* (Sheldrick 1997) and *WORDPERFECT* macro *PRPKAPPA* (Ferguson, 1999).

The authors thank ‘Servicios Técnicos de Investigación of Universidad de Jaén’ and the staff for the data collection, and Dr Manuel Melguizo for his help and advice. Lígia Gomes and Luís Santos acknowledge financial support from Fundação para Ciência e Tecnologia (POCI/QUI/61873/2004).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2646).

References

- Crispini, A. & Neve, F. (2002). *Acta Cryst. C58*, o34–o35.
- Duisenberg, A. J. M., Hooft, R. W. W., Schreurs, A. M. M. & Kroon, J. (2000). *J. Appl. Cryst. 33*, 893–898.
- Duisenberg, A. J. M., Kroon-Batenburg, L. M. J. & Schreurs, A. M. M. (2003). *J. Appl. Cryst. 36*, 220–229.
- Ferguson, G. (1999). *PRPKAPPA*. University of Guelph, Canada.
- Miyaura, N. & Suzuki, A. (1995). *Chem. Rev. 95*, 2457–2483.
- Nonius (1999). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Oszlányi, G. & Sütő, A. (2004). *Acta Cryst. A60*, 134–141.
- Oszlányi, G. & Sütő, A. (2005). *Acta Cryst. A61*, 147–152.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Sheldrick, G. M. (2003). *SADABS*. Version 2.10. Bruker AXS Inc., Madison, Wisconsin, USA.
- Silva, A. M. S., Almeida, L. M. P. M., Cavaleiro, J. A. S., Foces-Foces, C., Liamas-Saiz, A. L., Fontenás, C., Jagerovic, N. & Elguero, J. (1997). *Tetrahedron*, **53**, 11645–11658.
- Spek, A. L. (2003). *J. Appl. Cryst. 36*, 7–13.

supplementary materials

Acta Cryst. (2007). E63, o4833 [doi:10.1107/S1600536807058850]

2,6-Diphenylpyridine

M. A. Rocha, J. N. Low, L. R. Gomes, A. Quesada and L. M. N. B. F. Santos

Comment

The bonds and angles are in agreement with those found for related structures described by Crispini & Neve. (2002) and Silva *et al.* (1997). The angles between the mean planes of the pyridine rings and the phenyl rings are 29.68 (18) $^{\circ}$ (ring attached to C2), and 26.58 (17) $^{\circ}$ (ring attached to C6). The molecules are linked into chains along [0 $\bar{1}$ 1] by a weak C—H.. π interaction (Fig. 2 and Table 1); C64—H64· $Cg1(-x + 1/2, y + 1/2, z - 1/2)$, where $Cg1$ is the centroid of the pyridine ring. There are no further intermolecular interactions.

Experimental

The title compound was synthesized by adapting the procedure described by Miyaura & Suzuki (1995). A solution of 2,6-dibromopyridine (1.9 g), palladium(II) acetate (0.02 g) and phenylboronic acid (5.5 g) in *N,N*'-dimethylformamide was added to an aqueous solution of potassium carbonate (4.3 g). The mixture was refluxed for 5 h at 373 K. The cooled solution was extracted with 4 \times 50 ml ethyl acetate/acetone (5:1 v/v) and the extract was filtered out. The solvent was evaporated under low pressure at 353 K. The resulting solid was recrystallized from ethanol to yield colourless blocks of (I). Found (%w) for C₁₇H₁₃N: C 88.97, H 5.32, N 5.71; calculated (%w): C 88.28, H 5.67, N 6.06. m.p. 351 K.

Refinement

Anomalous dispersion was negligible and Friedel pairs were merged before refinement. H atoms were treated as riding atoms with C—H = 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. A view of (I) with displacement ellipsoids drawn at the 30% probability level (arbitrary spheres for the H atoms).




Fig. 2. A stereoscopic view of symmetry related chains of molecules linked by the C—H.. π interaction. H atoms not involved in the interaction are omitted for clarity.

supplementary materials

2,6-Diphenylpyridine

Crystal data

C ₁₇ H ₁₃ N	D _x = 1.206 Mg m ⁻³
M _r = 231.28	Mo K α radiation
Orthorhombic, Pna2 ₁	λ = 0.71073 Å
a = 16.1368 (16) Å	Cell parameters from 1596 reflections
b = 12.5371 (14) Å	θ = 3.5–27.5°
c = 6.2969 (4) Å	μ = 0.07 mm ⁻¹
V = 1273.9 (2) Å ³	T = 120 (2) K
Z = 4	Block, colourless
F ₀₀₀ = 488	0.70 × 0.32 × 0.14 mm

Data collection

Bruker-Nonius CCD diffractometer	1596 independent reflections
Radiation source: Bruker-Nonius FR591 rotating anode	1147 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.053$
Detector resolution: 9.091 pixels mm ⁻¹	$\theta_{\text{max}} = 27.5^\circ$
T = 120(2) K	$\theta_{\text{min}} = 3.5^\circ$
π & ω scans	$h = -20 \rightarrow 20$
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)	$k = -16 \rightarrow 16$
$T_{\text{min}} = 0.953$, $T_{\text{max}} = 0.990$	$l = -8 \rightarrow 8$
29987 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.055$	H-atom parameters constrained
$wR(F^2) = 0.163$	$w = 1/[\sigma^2(F_o^2) + (0.0775P)^2 + 0.6523P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\text{max}} < 0.001$
1596 reflections	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
163 parameters	$\Delta\rho_{\text{min}} = -0.25 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

supplementary materials

Special details

Experimental. The scale factors in the experimental table are calculated from the 'size' command in the *SHELXL97* input file.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.40272 (15)	0.4122 (2)	0.3621 (5)	0.0353 (6)
C2	0.4510 (2)	0.3407 (3)	0.4621 (6)	0.0356 (8)
C3	0.4702 (2)	0.3502 (3)	0.6756 (6)	0.0412 (8)
C4	0.4376 (2)	0.4338 (3)	0.7908 (7)	0.0443 (9)
C5	0.3893 (2)	0.5073 (3)	0.6890 (6)	0.0406 (8)
C6	0.37309 (19)	0.4956 (2)	0.4725 (6)	0.0345 (7)
C21	0.4814 (2)	0.2503 (3)	0.3325 (6)	0.0407 (8)
C22	0.5556 (2)	0.1994 (3)	0.3788 (8)	0.0503 (10)
C23	0.5826 (3)	0.1144 (3)	0.2567 (9)	0.0624 (13)
C24	0.5356 (3)	0.0782 (3)	0.0889 (9)	0.0635 (12)
C25	0.4618 (3)	0.1281 (3)	0.0410 (8)	0.0594 (12)
C26	0.4351 (3)	0.2137 (3)	0.1592 (6)	0.0468 (9)
C61	0.32310 (19)	0.5744 (3)	0.3538 (6)	0.0386 (8)
C62	0.3186 (2)	0.6797 (3)	0.4181 (7)	0.0512 (10)
C63	0.2743 (3)	0.7541 (3)	0.3035 (9)	0.0613 (12)
C64	0.2330 (2)	0.7237 (4)	0.1205 (9)	0.0629 (13)
C65	0.2361 (2)	0.6181 (3)	0.0548 (8)	0.0556 (11)
C66	0.2814 (2)	0.5446 (3)	0.1690 (6)	0.0434 (9)
H3	0.5055	0.2996	0.7421	0.049*
H4	0.4486	0.4402	0.9385	0.053*
H5	0.3668	0.5659	0.7652	0.049*
H22	0.5881	0.2231	0.4953	0.060*
H23	0.6338	0.0808	0.2888	0.075*
H24	0.5539	0.0192	0.0067	0.076*
H25	0.4292	0.1032	-0.0744	0.071*
H26	0.3846	0.2483	0.1229	0.056*
H62	0.3466	0.7013	0.5437	0.061*
H63	0.2721	0.8261	0.3500	0.074*
H64	0.2028	0.7748	0.0405	0.076*
H65	0.2069	0.5965	-0.0691	0.067*
H66	0.2842	0.4728	0.1214	0.052*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0342 (13)	0.0336 (14)	0.0381 (15)	-0.0040 (11)	0.0018 (14)	0.0020 (13)
C2	0.0335 (16)	0.0326 (16)	0.0408 (19)	-0.0040 (13)	0.0022 (15)	0.0032 (16)
C3	0.0344 (16)	0.049 (2)	0.041 (2)	-0.0077 (15)	-0.0038 (16)	0.0062 (18)
C4	0.0417 (18)	0.052 (2)	0.039 (2)	-0.0110 (16)	-0.0035 (17)	0.0011 (18)
C5	0.0391 (17)	0.0454 (19)	0.0371 (19)	-0.0072 (15)	0.0058 (16)	-0.0080 (17)

supplementary materials

C6	0.0311 (15)	0.0338 (17)	0.0384 (19)	-0.0021 (13)	0.0009 (15)	-0.0015 (16)
C21	0.0453 (18)	0.0320 (15)	0.045 (2)	0.0002 (14)	0.0075 (18)	0.0058 (16)
C22	0.049 (2)	0.0365 (18)	0.066 (3)	0.0026 (15)	0.006 (2)	0.012 (2)
C23	0.063 (2)	0.039 (2)	0.085 (3)	0.0123 (18)	0.019 (3)	0.013 (2)
C24	0.090 (3)	0.0358 (18)	0.065 (3)	0.011 (2)	0.024 (3)	0.003 (2)
C25	0.088 (3)	0.0359 (19)	0.054 (3)	0.005 (2)	0.006 (3)	0.0004 (19)
C26	0.061 (2)	0.0344 (18)	0.045 (2)	0.0022 (16)	0.000 (2)	0.0007 (17)
C61	0.0307 (14)	0.0407 (17)	0.0444 (19)	0.0019 (13)	0.0034 (17)	-0.0021 (16)
C62	0.050 (2)	0.045 (2)	0.059 (3)	0.0085 (17)	0.0038 (19)	-0.005 (2)
C63	0.056 (2)	0.048 (2)	0.079 (4)	0.0154 (18)	0.011 (2)	0.000 (2)
C64	0.0384 (19)	0.068 (3)	0.083 (4)	0.0231 (19)	0.003 (2)	0.017 (3)
C65	0.0374 (19)	0.073 (3)	0.057 (3)	0.0057 (19)	-0.0049 (19)	0.005 (2)
C66	0.0331 (16)	0.0495 (19)	0.048 (2)	0.0032 (15)	0.0001 (18)	0.0013 (19)

Geometric parameters (\AA , °)

N1—C6	1.344 (4)	C24—C25	1.378 (7)
N1—C2	1.344 (4)	C24—H24	0.95
C2—C3	1.385 (5)	C25—C26	1.376 (5)
C2—C21	1.481 (5)	C25—H25	0.95
C3—C4	1.378 (5)	C26—H26	0.95
C3—H3	0.95	C61—C62	1.383 (5)
C4—C5	1.368 (5)	C61—C66	1.395 (5)
C4—H4	0.95	C62—C63	1.379 (6)
C5—C6	1.395 (5)	C62—H62	0.95
C5—H5	0.95	C63—C64	1.384 (7)
C6—C61	1.478 (5)	C63—H63	0.95
C21—C22	1.387 (5)	C64—C65	1.388 (6)
C21—C26	1.400 (5)	C64—H64	0.95
C22—C23	1.384 (6)	C65—C66	1.380 (5)
C22—H22	0.95	C65—H65	0.95
C23—C24	1.378 (8)	C66—H66	0.95
C23—H23	0.95		
C6—N1—C2	118.9 (3)	C25—C24—H24	120.2
N1—C2—C3	121.8 (3)	C23—C24—H24	120.2
N1—C2—C21	116.4 (3)	C26—C25—C24	120.5 (5)
C3—C2—C21	121.8 (3)	C26—C25—H25	119.8
C4—C3—C2	119.4 (4)	C24—C25—H25	119.8
C4—C3—H3	120.3	C25—C26—C21	120.6 (4)
C2—C3—H3	120.3	C25—C26—H26	119.7
C5—C4—C3	118.9 (4)	C21—C26—H26	119.7
C5—C4—H4	120.6	C62—C61—C66	118.3 (3)
C3—C4—H4	120.6	C62—C61—C6	121.2 (3)
C4—C5—C6	119.6 (3)	C66—C61—C6	120.4 (3)
C4—C5—H5	120.2	C63—C62—C61	121.3 (4)
C6—C5—H5	120.2	C63—C62—H62	119.3
N1—C6—C5	121.3 (3)	C61—C62—H62	119.3
N1—C6—C61	116.9 (3)	C62—C63—C64	119.9 (4)
C5—C6—C61	121.7 (3)	C62—C63—H63	120.0

supplementary materials

C22—C21—C26	118.3 (4)	C64—C63—H63	120.0
C22—C21—C2	121.5 (4)	C63—C64—C65	119.6 (4)
C26—C21—C2	120.2 (3)	C63—C64—H64	120.2
C23—C22—C21	120.6 (4)	C65—C64—H64	120.2
C23—C22—H22	119.7	C66—C65—C64	120.0 (4)
C21—C22—H22	119.7	C66—C65—H65	120.0
C24—C23—C22	120.4 (4)	C64—C65—H65	120.0
C24—C23—H23	119.8	C65—C66—C61	120.8 (4)
C22—C23—H23	119.8	C65—C66—H66	119.6
C25—C24—C23	119.6 (4)	C61—C66—H66	119.6
C6—N1—C2—C3	0.7 (5)	C22—C23—C24—C25	0.9 (7)
C6—N1—C2—C21	179.7 (3)	C23—C24—C25—C26	0.1 (7)
N1—C2—C3—C4	1.4 (5)	C24—C25—C26—C21	-1.2 (6)
C21—C2—C3—C4	-177.5 (3)	C22—C21—C26—C25	1.3 (5)
C2—C3—C4—C5	-2.1 (5)	C2—C21—C26—C25	-178.4 (4)
C3—C4—C5—C6	0.8 (5)	N1—C6—C61—C62	-152.9 (3)
C2—N1—C6—C5	-2.1 (4)	C5—C6—C61—C62	26.8 (5)
C2—N1—C6—C61	177.6 (3)	N1—C6—C61—C66	25.0 (4)
C4—C5—C6—N1	1.4 (5)	C5—C6—C61—C66	-155.3 (3)
C4—C5—C6—C61	-178.3 (3)	C66—C61—C62—C63	-0.1 (6)
N1—C2—C21—C22	151.8 (3)	C6—C61—C62—C63	177.7 (4)
C3—C2—C21—C22	-29.2 (5)	C61—C62—C63—C64	0.2 (6)
N1—C2—C21—C26	-28.5 (5)	C62—C63—C64—C65	0.5 (6)
C3—C2—C21—C26	150.5 (3)	C63—C64—C65—C66	-1.3 (6)
C26—C21—C22—C23	-0.2 (5)	C64—C65—C66—C61	1.4 (6)
C2—C21—C22—C23	179.4 (4)	C62—C61—C66—C65	-0.7 (5)
C21—C22—C23—C24	-0.8 (6)	C6—C61—C66—C65	-178.6 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C64—H64 \cdots Cg1 ⁱ	0.95	2.74	3.534 (4)	142

Symmetry codes: (i) $-x+1/2, y+1/2, z-1/2$.

supplementary materials

Fig. 1




Fig. 2

